

WST-1 Cell Viability & Proliferation Assay

Cat. No. CB018

(1000 Tests in 96-well plate)

Product Description

The reduction of tetrazolium salts to colored formazan compounds by succinate-tetrazolium reductase, which exists in viable cells, provides a sensitive and accurate method to measure cell viability and proliferation. The most commonly used tetrazolium salt, MTT [3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide], however, suffers from the disadvantage that the formazan dye produced from MTT is extremely water insoluble, so an additional extraction step is needed for spectrophotometric quantification. Instead of MTT, the ScienCell™ WST-1 Cell Viability & Proliferation Assay utilizes a tetrazolium salt WST-1[2-(4-Iodophenyl)-3-(4-nitrophenyl)-5-(2,4-disulfophenyl)-2H-tetrazolium]. WST-1 produces a highly water soluble formazan upon metabolically active cells, allowing a direct and user-friendly colorimetric measurement of cell viability and proliferation.

Kit Components

WST-1 powder, 6.52 mg

Electro Coupling Reagent, 2 ml

Procedures (96-well plate)

1. Plate and culture cells in a clear-bottom 96-well tissue culture plate. Incubate cells with test compounds and controls for the desired period of time. The final volume of culture medium in each well should be 100 μ l.
2. Thaw Electro Coupling Reagent, and reconstitute each vial of WST-1 with 2 ml of Electro Coupling Reagent. Vortex briefly and keep in the dark at 4°C until use. Freshly reconstituted WST-1 is recommended for each experiment. For longer storage, we suggest that you aliquot and store the reconstituted WST-1 reagent at -20°C, avoid repeated freeze/thaw cycles.
3. Add 10 μ l of reconstituted WST-1 reagent to each well of 96-well plate (the volume of the WST-1 reagent should be 1/10 of the original culture medium). Mix well by gently rocking the plate side-to-side.
4. Incubate cultures with WST-1 at 37°C for 2-4 hours depending on cell type and seeding density.
5. After incubation, measure the absorbance on an ELISA plate reader with a test wavelength at 450 nm and a reference wavelength at 630 nm, and subtract the 630 nm background absorbance from the 450 nm measurement.

Usage

MTT Cell Viability & Proliferation Assay is used to coat cell culture vessels *in vitro*. MTT Cell Viability & Proliferation Assay is for research use only. It is not approved for human or animal use, or application in clinical or *in vitro* diagnostic procedures.